

Institute and Faculty of Actuaries

Reimagining the 1918 Pandemic

Ashley Campbell RMS LifeRisks

February 2018

Agenda

1. Influenza Characteristics

2. The 1918 Pandemic

3. Reimagining the 1918 Pandemic




Influenza Characteristics




Types of Influenza

- Type A
 - Zoonotic
 - Drifts frequently
 - Shifts cause pandemics
 - Group 1 & Group 2 strains



- Type B Human-to-Human & Seals; Not associated with pandemics
- Type C Humans & Pigs; Not associated with pandemics

Two phylogenetic groups of influenza



... e.g. H1N1


Contagion characteristics of an influenza virus

• Human-to-human contact





• Virus attaches to upper respiratory tract

Social Network is crucial for determining contagion



Shifts Cause Pandemics





H16 н13

H7

Group 2

H15 📩

H11

H9 •H12 **H**8

•H6


Most recent influenza pandemics

February 2018



The1918 Spanish Influenza – What happened?





The influenza pandemic of 1918 to 1919 is the deadliest in modern history.

~1/3 of world infected ~50 million deaths

High excess mortality in young people

Origin

• The 1918 influenza pandemic is mistakenly called 'The <u>Spanish</u> Flu'


 In late 1917, a virulent respiratory disease of unknown origins erupted in the interior of northern China

 China mobilized 95,000 labourers to support the war effort






Demographic response – Excess Mortality




From Taubenberger & Morens (2006): **1918** Influenza: the Mother of All Pandemics

Antigenic imprinting




- Child imprints on the HA group of their first flu exposure
- Reduced risk of severe disease from flu strains within that same phylogenetic group
- More vulnerable to other HA group strains

Antigenic imprinting: 1918 Pandemic




Antigenic Imprinting: 2009 "Swine-Flu"




• 151,700 – 575,400 respiratory pandemic deaths Globally

Antigenic Imprinting: 2009 "Swine-Flu"


2009 Pandemic age shift Pneumonia mortality in Mexico



Antigenic imprinting: "Bird-Flu" Epidemic



Antigenic imprinting: "Bird-Flu" Epidemic






Reimagining the 1918 pandemic




Scenario Structure


- Transmissibility = 2
- Lethality = 2.5%
- Assume vaccine available after 6 months
- Assume ineffective Pharmaceuticals




RMS Infectious Disease Model Structure

Demographic profile






February 2018




February 2018

Results


- Most exposures were born after 1968. Vulnerable to group 1





Conclusions

- Cohort effects exist and can change over time depending on circulating flu strains at birth



- Most life insurers will be more vulnerable to Group 1, e.g. H1, H2, H5, H6



Expressions of individual views by members of the Institute and Faculty of Actuaries and its staff are encouraged.

The views expressed in this presentation are those of the presenter.