



Actuarial Research Centre

Institute and Faculty
of Actuaries

Actuarial Research Centre (ARC)

PhD studentship output

The Actuarial Research Centre (ARC) is the Institute and Faculty of Actuaries' network of actuarial researchers around the world. The ARC seeks to deliver research programmes that bridge academic rigour with practitioner needs by working collaboratively with academics, industry and other actuarial bodies.

The ARC supports actuarial researchers around the world in the delivery of cutting-edge research programmes that aim to address some of the significant challenges in actuarial science.



Actuarial
Research Centre

Institute and Faculty
of Actuaries

Bayesian Inference for Small Population Longevity Risk Modelling

Liang Chen, CERA

Supervised by: Prof Andrew Cairns, Dr Torsten Kleinow

Actuarial Research Centre, IFoA
Heriot Watt University

July 25, 2016

Stochastic Model

We select stochastic model “M7” to reflect the work of Cairns et al. (2009), which suggests it fit the males from England and Wales well.

Recall the formula for M7:

$$D(t, x) | \theta_1 \sim Poi(m(\theta_1, t, x)E(t, x))$$

$$\text{logit } q(\theta_1, t, x) = \kappa_t^{(1)} + \kappa_t^{(2)}(x - \bar{x}) + \kappa_t^{(3)}((x - \bar{x})^2 - \hat{\sigma}_x^2) + \gamma_c^{(4)}$$

- $\theta_1 = (\kappa_t^{(1)}, \kappa_t^{(2)}, \kappa_t^{(3)}, \gamma_c^{(4)})$
- $\kappa_t^{(i)}$ is a period effect in year $t = t_1, \dots, t_{n_y}$ for each $i = 1, 2, 3$.
- $\gamma_c^{(4)}$ is the cohort effect for the cohort born in year $c = t - x$ for $t = t_1, \dots, t_{n_y}$ and $x = x_1, \dots, x_{n_a}$.
- \bar{x} is the mean of the age range we use for our analysis.
- $\hat{\sigma}_x^2$ is the mean of $(x - \bar{x})^2$.



Two-Stage Approach

Stage

1. Find the estimates for period and cohort effects, $\hat{\theta}_1$ by maximising the Poisson likelihood.
2. Fit time series model to these effects.

Most pension schemes are less than 1% of national population.

Two-stage approach leads to biased estimates of volatility for small populations.

- Large sampling variation affects latent parameter estimation, with significant noise obscuring the true signal (Cairns, Blake, Dowd et al. 2011).
- Results in non-negligible bias to the parameter estimation of the projecting model, given the assumed true rates (Chen, Cairns and Kleinow 2015).
- Over fit the cohorts with only one observation (a problem with the two-stage approach: see Cairns et al. 2009)



Bayesian Approach

Bayesian approach offers a way to avoid or reduce this bias by

- Combining Poisson and time series likelihoods
- Using knowledge of larger England and Wales dataset to choose more informative priors than one might normally choose.

We use England and Wales death rates as a benchmark to test how well the Bayesian approach with informative priors performs.



**Actuarial
Research Centre**

Institute and Faculty
of Actuaries

Data

- Benchmark exposure $E_0(t, x)$ and corresponding deaths count $D_0(t, x)$ of the males in England and Wales (EW) in the HMD database, during year 1961 to 2011, aged 50-89 last birthday.
- Simulate $D_w(t, x)$, where $w = 0.01$ based on
$$D_w(t, x) | \hat{\theta}_0 \sim \text{Poi}(m(\hat{\theta}_0, t, x)wE_0(t, x))$$

where

- $\hat{\theta}_0$: parameter estimates for benchmark $D_0(t, x)$, i.e. EW
- $m(\hat{\theta}_0, t, x)$ is the fitted death rates given $\hat{\theta}_0$, that is $\hat{\theta}_0$ is the true rates for $D_w(t, x)$.
- Find the parameter estimates $\hat{\theta}_w$ for $D_w(t, x)$.



Actuarial
Research Centre

Institute and Faculty
of Actuaries

Notations

- θ_1 , the vector of all the latent parameters
- $\theta_{11} = \left(\kappa_{t_1}^{(1)}, \kappa_{t_1}^{(2)}, \kappa_{t_1}^{(3)} \right)^T$, vector of period effects at year t_1
- θ_{12} , vector of the rest of period effects
- $\theta_{13} = \gamma_{t_1 - x_{n_a}}^{(4)}$
- θ_{14} , vector of cohort effect for the rest cohorts



Prior for κ and $\gamma^{(4)}$

- $\theta_{11} \propto$ uniform distribution
- θ_{12} , multivariate random walk:

$$\kappa_t = \kappa_{t-1} + \mu + \epsilon_t$$

where

- $\mu = (\mu_1, \mu_2, \mu_3)^T$ is the drift (hyper-parameter).
- $\epsilon_t \sim MVN(\mathbf{0}, \mathbf{V}_\epsilon)$, *i.i.d* three dimensional multivariate normal distribution independent of t .
- θ_{14} , AR(1) model:

$$\gamma_c^{(4)} = \alpha_\gamma \gamma_{c-1}^{(4)} + \epsilon_c, \text{ for } c > t_1 - x_{n_a},$$

where ϵ_c are *i.i.d* and $\epsilon_c \sim N(0, \sigma_\gamma^2)$.

- $\gamma_c^{(4)} | \gamma_{c-1}^{(4)} \sim N(\alpha_\gamma \gamma_{c-1}^{(4)}, \sigma_\gamma^2)$
- $\gamma_{t_1}^{(4)} \sim N(0, \frac{\sigma_\gamma^2}{1-\alpha_\gamma^2})$




Prior for Hyper-Parameters





- $V_\epsilon \propto$ Inverse Wishart (ν, Σ)
 - MCMC-Mean: Fix the mean of prior to $\widehat{V}_\epsilon^{EW}$
 - MCMC-Mode: Fix the mode of prior to $\widehat{V}_\epsilon^{EW}$ (sensitivity test)
- $\mu \propto$ uniform
- $\alpha_\gamma \propto (1 - \alpha_\gamma^2)^g$ for $|\alpha| < 1$
- $\sigma_\gamma^2 \sim$ Inverse Gamma (a_γ, b_γ)




V_ϵ given MLE




Credibility Interval for κ and $\gamma^{(4)}$




CDF for μ_1 with Sensitivity Test




CDF for $V_{\epsilon}(1, 1)$ with Sensitivity Test




CDF for μ_2 with Sensitivity Test




CDF for $V_{\epsilon}(2, 2)$ with Sensitivity Test



CDF for μ_3 with Sensitivity Test



CDF for $V_{\epsilon}(3, 3)$ with Sensitivity Test



Conclusion

For small population

- The co-variance matrix estimated by MLE is significantly biased to the right of the assumed true value due to the Poisson model's over fitting.
- We combined the two stages into one by adding time series likelihood for the latent parameters and gained the posterior distribution with the MCMC procedure.
- The Bayesian method provides an improved fit to the hyper parameter V_{ϵ} .
- The low level information involved in short cohorts is balanced by the time series prior.
- The posterior distribution for small population is sensitive and fixing the mode of the prior for the co-variance matrix to the assumed true rates provides approximately unbiased fit to V_{ϵ}



Questions

Comments



**Actuarial
Research Centre**

Institute and Faculty
of Actuaries