THE RECENT TREND OF MORTALITY IN ENGLAND AND WALES

By W. S. HOCKING, O.B.E., F.I.A. of the Government Actuary's Department

In continuation of the tables given in J.I.A. Vol. LXXIII, pp. 431-34, the central death-rates in age-groups for males and females separately in the year 1946 are shown in Table 1, together with the ratios of these death-rates to the corresponding rates for the years 1930-31-32. Both the deaths and the populations from which the death-rates have been derived are limited to civilians, but civilian deaths due to war causes are included. The effect of the withdrawal of healthy lives to the Forces should be borne in mind in examining the rates for young adult ages.

The tables showing the death-rates and the ratios to the 1930–32 standard are scattered over a number of volumes of the Journal. In order that the trend up to this first post-war year may be conveniently studied, two new tables have been included on this occasion, showing the rates (Table 2) and ratios (Table 3) for each year since 1933. In these tables the age-groups have been reduced to six, viz. under age 1 (infants), 1 to 4 (young children), 5 to 14 (children of school age), 15 to 44 (younger adults), 45 to 64 (older adults) and 65 and over (elderly). Further, to eliminate the effect of the considerable changes in the age distribution of the population that have taken place since 1930–32, the group death-rates and ratios shown in these tables have been standardized on the age distribution of England and Wales at the 1931 census. For the benefit of those who prefer charts, the ratios to the 1930–32 standard have also been shown in graphical form.

Table 1. Death-rates for age-groups in the year 1946 in England and Wales*

| Age-group | Death | ı-rates | Ratios of death-rates to those for 1930-31-32 | | |
|-----------|-----------------|---------|---|---------|--|
| | Males | Females | Males | Females | |
| 0-4 | .01343 | .01036 | .626 | .615 | |
| 5-9 | .00097 | .00074 | .424 | ·365 | |
| 10-14 | 00084 | .00065 | .560 | 451 | |
| 15-19 | .00159 | .00110 | .624 | .517 | |
| 20-24 | .00271 | .00180 | ·826 | 641 | |
| 25-29 | .00201 | .00194 | .609 | .634 | |
| 30-34 | .00227 | .00206 | .624 | -617 | |
| 35-39 | .00283 | '00241 | .593 | .612 | |
| 40-44 | .00414 | .00312 | .642 | .643 | |
| 45-49 | 100691 | .00485 | 742 | .722 | |
| 50-54 | ·01094 | .00679 | ·836 | .716 | |
| 55-59 | 01718 | .01002 | ∙899 | .723 | |
| 60-64 | .02741 | .01624 | .939 | .761 | |
| 65-69 | .04012 | .02583 | -863 | .765 | |
| 70-74 | .06311 | .04576 | ·848 | 824 | |
| 75-79 | · 0 9978 | .07609 | ·847 | 818 | |
| 80-84 | 15951 | 12233 | ∙888 | ·828 | |
| 85 & over | .21734 | -20286 | .771 | .819 | |
| All ages† | .00999 | .00822 | ·786 | .731 | |

^{*} See note on p. 346.

[†] Standardized on 1931 census age distribution.

Table 2. Standardized death-rates for age-groups in England and Wales (standardized on 1931 census age distribution)

| | Age-group | | | | | All | | |
|---------|------------|--------|--------|--------|--------|--------------|--------|--|
| Year | Under 1 | 1-4 | 5-14 | 15-44 | 45-64 | 65 & over | ages | |
| | Males | | | | | | | |
| 1930-32 | .07650 | .00755 | -00190 | .00385 | .01653 | .08063 | .01271 | |
| 1933 | .07354 | ·00688 | .00188 | .00390 | ·01687 | .08082 | .01273 | |
| 1934 | .07109 | .00697 | .00197 | .00356 | .01628 | .07702 | 01217 | |
| 1935 | ·06791 | .00535 | .00174 | .00342 | .01911 | .07775 | .01192 | |
| 1936 | .07042 | -00580 | .00170 | .00332 | .01655 | .08037 | .01221 | |
| 1937 | ·o6845 | .00536 | .00160 | -00338 | ·01698 | ·08131 | 01231 | |
| 1938 | .06257 | .00483 | .00160 | .00319 | .01569 | .07618 | .01146 | |
| 1939 | .05874 | .00367 | .00137 | .00304 | .01628 | 07947 | .01128 | |
| 1940 | .06631 | .00211 | 18100. | .00447 | .01923 | .08729 | .01368 | |
| 1941 | .07111 | .00557 | .00188 | .00479 | .01201 | 07774 | .01285 | |
| 1942 | .06150 | .00372 | .00142 | .00393 | .01203 | 07089 | .01122 | |
| 1943 | .05746 | ·00348 | .00136 | .00407 | '01542 | .07146 | '01132 | |
| 1944 | .05428 | .00298 | .00138 | .00409 | .01517 | ·06891 | .01105 | |
| 1945 | 05194 | .00275 | .00110 | .00385 | .01477 | ·06849 | .01071 | |
| 1946* | .05559 | .00221 | .00001 | .00251 | .01443 | ·06861 | .00999 | |
| | Females | | | | | | | |
| 1930-32 | .05705 | .00677 | .00174 | '00333 | .01197 | .06839 | '01124 | |
| 1933 | .05587 | .00624 | .00176 | .00342 | .01215 | 07063 | .01145 | |
| 1934 | .05486 | .00632 | .00100 | .00314 | .01138 | .06537 | .01074 | |
| 1935 | .05207 | .00489 | .00165 | 100302 | .01118 | .06555 | .01048 | |
| 1936 | .05286 | .00218 | .00120 | .00290 | .01130 | .06793 | .0106 | |
| 1937 | .05235 | .00476 | .00144 | .00293 | .01145 | .06919 | .01076 | |
| 1938 | .04738 | .00423 | .00145 | .00264 | .01046 | .06273 | .00970 | |
| 1939 | .04566 | .00322 | .00114 | .00260 | .01071 | .06547 | .00992 | |
| 1940 | .05000 | ·00444 | .00156 | .00318 | .01208 | .07194 | .0112 | |
| 1941 | .05462 | .00499 | .00156 | .00302 | .01062 | .06323 | .0102 | |
| 1942 | 04752 | .00318 | .00111 | .00254 | .00921 | .05633 | .00890 | |
| 1943 | 04478 | .00312 | 80100 | 00256 | .00958 | .05847 | .0090 | |
| 1944 | .04254 | .00254 | .00102 | .00247 | .00921 | .05462 | .0085 | |
| 1945 | .04063 | .00232 | .00000 | .00224 | .00899 | .05485 | .00837 | |
| 1946* | '04229 | .00185 | .00070 | '00205 | .00879 | .05543 | .00822 | |


^{*} After these Tables had gone to press, the Registrar-General's estimate of the age distribution of the population was revised. This revision results in alterations of 2 per cent or more in a number of age groups, and for the age group 85 and over for men there is an increase in the mortality rate of as much as 7 per cent. Group Tables based on the revised estimate of the population will be included in the next number of the Journal.

Table 3. Ratios of standardized death-rates for age-groups in England and Wales to the rates for 1930-31-32 (standardized on 1931 census age distribution)

| *** | Age-group | | | | | | All |
|---------|---------------|-------|-------|-------|-------|-----------|-------|
| Year - | Under 1 | 1-4 | 5-14 | 15-44 | 45-64 | 65 & over | ages |
| | Males | | | | | | |
| 1930-32 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1,000 | 1.000 |
| 1933 | ·961 | .911 | -989 | 1.013 | 1.021 | 1.003 | 1.002 |
| 1934 | .929 | .923 | 1.037 | ·925 | .985 | ·955 | .958 |
| 1935 | ⋅888 | .709 | ·916 | ·888 | .975 | ·964 | .938 |
| 1936 | .921 | .768 | 895 | ·862 | 1.001 | .997 | .961 |
| 1937 | ·895 | .710 | ·842 | ⋅878 | 1.027 | 1.008 | .969 |
| 1938 | -818 | -640 | 842 | ·821 | .949 | ·945 | .902 |
| 1939 | ·768 | ·486 | .721 | •790 | .985 | ·986 | 116. |
| 1940 | ·867 | •677 | .953 | 1.161 | 1.163 | 1.083 | 1.076 |
| 1941 | .930 | .738 | •989 | 1.244 | 1.020 | .964 | 1.011 |
| 1942 | ·8 0 4 | .493 | .747 | 1.021 | .909 | .879 | .883 |
| 1943 | 751 | -461 | -716 | 1.057 | -933 | ∙886 | .891 |
| 1944 | .710 | .395 | .726 | 1.063 | .918 | ·855 | .867 |
| 1945 | •679 | ٠364 | ·626 | 1.000 | ٠894 | ·849 | ·843 |
| 1946* | .727 | .293 | .479 | .652 | ·873 | .851 | .786 |
| | Females | | | | | | |
| 1930-32 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 | 1.000 |
| 1933 | .979 | .922 | 1.011 | 1.027 | 1.012 | 1.033 | 1.010 |
| 1934 | .962 | .934 | 1.002 | .943 | .951 | ·956 | ·956 |
| 1935 | .913 | .722 | .931 | 907 | .934 | .958 | .932 |
| 1936 | .927 | .765 | .862 | ·871 | .944 | .993 | .948 |
| 1937 | 918 | .703 | ·828 | ·88o | .957 | 1.013 | .957 |
| 1938 | .831 | ·625 | .833 | 793 | ·874 | .917 | ·871 |
| 1939 | -800 | .476 | .655 | ·781 | .895 | .957 | -883 |
| 1940 | ·876 | 656 | ·897 | :955 | 1.000 | 1.052 | .997 |
| 1941 | 957 | .737 | .897 | .907 | ·89î | ·925 | 908 |
| 1942 | ·833 | .470 | ·638 | .763 | .794 | .824 | .792 |
| 1943 | .785 | ·468 | .621 | .769 | -800 | .855 | 805 |
| 1944 | .746 | .375 | •603 | .742 | •769 | ·799 | ·76ŏ |
| 1945 | .712 | 347 | .517 | .673 | .751 | 1802 | .742 |
| 1946* | .741 | .273 | .402 | ·616 | .734 | ∙810 | ·731 |


^{*} See note on p. 346.

Ratios of group death-rates (standardized) to 1930-32 rates



Ratios markedly affected by war conditions have been joined together by broken lines.

Ratios of group death-rates (standardized) to 1930-32 rates



Ratios markedly affected by war conditions have been joined together by broken lines.