THE RECENT TREND OF MORTALITY IN GREAT BRITAIN

By L. V. MARTIN, F.I.A.

of the Government Actuary's Department

In continuation of the tables given in J.I.A. 87, 384-88, central deathrates in age-groups for males and females separately in the year 1961, and the ratios of these death-rates to the corresponding rates for the years 1950-52 taken as the standard, are shown in Table 1. The deaths

Table 1. Death-rates for age-groups in Great Britain in the year 1961 and ratios of these rates to the rates for 1950–52

Death-rat	es in 1961	Ratios to 1950-52 rates		
Males	Females	Males	Females	
·02546	·01976	.74	.75	
.00106	·00082	·7 5	•68	
.00049	.00033	.72	•69	
	.00025	•71	•64	
	.00039	•95	.57	
.00111	.00021	∙80	.52	
.00106	.00058	.70	.47	
.00129	-00089	·74	·61	
.00194	.00138	⋅83	.72	
.00308	.00224	·8 ₇	∙8₃	
.00537	.00355	∙85	·8 ₄	
.00951	.00555	∙86	∙86	
.01716	·00834	•94	·8 ₄	
.02906	.01400	•99	∙88	
.0446	0235	•99	∙88	
	.0414	1.00	.90	
-	.0700	.95	∙88	
	.120	∙96	∙89	
.257	.228	•90	·97	
.01263	01139	.992	1.022	
·01186	•00969	.932	•869	
	Males ·02546 ·00106 ·00049 ·00039 ·00011 ·00106 ·00129 ·00194 ·00308 ·00537 ·00951 ·01716 ·02906 ·0446 ·165 ·257	.02546 .01976 .00106 .00082 .00049 .00033 .00039 .00025 .00091 .00039 .00111 .00051 .00106 .00058 .00129 .00089 .00194 .00138 .00308 .00224 .00537 .00355 .00951 .00555 .01716 .00834 .02906 .01400 .0446 .0235 .0690 .0414 .1046 .0700 .165 .120 .257 .228	Males Females Males .02546 .01976 .74 .00106 .00082 .75 .00049 .00033 .72 .00039 .00025 .71 .00091 .00039 .95 .00111 .00051 .80 .00106 .00058 .70 .00129 .00089 .74 .00194 .00138 .83 .00308 .00224 .87 .00537 .00355 .85 .01716 .00834 .94 .02906 .01400 .99 .0446 .0235 .99 .0466 .0700 .95 .105 .120 .96 .257 .228 .90	

^{*} Standardized on 1951 Census age distribution.

are those recorded as occurring in Great Britain in the year and the population used is the 'home' population, i.e. the number of people actually in the country at a stated time, as estimated by the Registrars General of England and Wales and of Scotland.

The estimated population for 1961 has been adjusted to take account of the preliminary results of the 1961 Census, but the adjustments are relatively small and do not materially affect comparisons with earlier years. Further revision may be necessary when the age distribution of the population enumerated at the Census becomes available.




Fig. 1. Ratios of group death-rates (standardized) to 1950-52 rates—children.

In order that the trend of mortality rates may be conveniently studied, the group death-rates and the ratios to the 1950-52 standard since 1951 are set out in Tables 2 and 3, and the ratios to the 1950-52 standard are shown in graphical form in Figs. 1 and 2. Figures for 1950 and

earlier years have not been given, but they are available in previous articles. In these tables and graphs the age-groups have been reduced to six under age I (infants), I-4 (young children), 5-I4 (children of school age), 15-44 (younger adults), 45-64 (older adults) and 65 and over (elderly).




Fig. 2. Ratios of group death-rates (standardized) to 1950-52 ratesadults and all ages.

Further, the group death-rates and ratios have been standardized on the age distribution of Great Britain at the 1951 Census so as to eliminate the effect of changes in the age distribution of the population.

As a link with the earlier series of tables that appeared in the *Journal*, showing the trend of mortality in England and Wales since 1930-32, figures for Great Britain for the average of those three years have been included in Tables 2 and 3.

Table 2. Standardized death-rates for age-groups in Great Britain (standardized on 1951 Census age distribution)

Age-group							
Year	Under 1	1-4	5-14	15-44	45-64	65 and over	All ages
			Ma	ales			
1930–32	· o 7953	.00760	.00196	.00413	.01637	.0873	.01288
1951	.03524	.00149	.00062	.00199	.01528	·0877	.01336
1952	.03293	.00134	.00022	·00184	·01404	· 0 787	.01211
1953	.03139	.00127	.00023	.00174	.01392	·0790	.01204
1954	.02992	.00108	·00046	·00170	·01382	·o787	01192
1955	02955	.00104	· ooo 49	.00162	·01382	·0807	.01208
1956	·02842	.00099	·00044	.00161	.01375	∙0804	.01199
1957	.02779	.00104	· ooo 48	·00164	.01391	.0774	·01176
1958	·02676	·00096	.00045	.00160	·01364	· 0 793	.01183
1959	· 02 597	.00100	·00044	.00191	·01358	·o784	.01173
196 0	· 02 663	·00094	· o oo46	.00160	.01335	· o 765	·01149
1961	·02546	.00103	.00044	.00160	.01363	· o 799	.01186
			Fem	nales			
1930-32	.05973	·oo686	.00178	.00321	.01243	.0735	.01510
1951	.02668	.00130	.00043	.00156	.00895	.0674	·01174
1952	·02556	.00106	.00039	·00136	.00822	· o 584	01032
1953	·02461	.00110	· oo o39	.00131	·00801	.0585	.01025
1954	.02272	•00083	.00034	.00127	.00794	.0572	.00999
1955	.02270	.00095	.00035	.00117	.00783	.0591	81010.
1956	.02153	.00082	.00031	.00112	.00770	.0585	.01002
1957	.02160	•00086	.00033	.00112	.00770	.0550	·00960
1958	·02068	.00075	.00028	.00109	.00755	.0567	00972
1959	.02062	.00081	.00031	.00107	.00742	.0561	.00961
1960	.02048	.00076	.00031	.00102	.00728	.0545	.00936
1961	· 0 1976	.00079	.00029	.00103	.00743	.0569	00969

The standardized death-rates in the year 1961 for all ages combined were heavier than in 1960 by about 3% for men and $3\frac{1}{2}$ % for women, but were 7% less for men and 13% less for women than the 1950–52 standard. At ages over 65 mortality was heavier than in any year since 1956, but at younger ages the rates differed little from those of the last few years and the infant mortality rates continue to decline.

Table 3. Ratios of standardized death-rates for age-groups in Great Britain to the rates for 1950–52 (standardized on 1951 Census age distribution)

Age-group							
Year	Under 1	1-4	5-14	15-44	45-64	65 and over	All ages
Males							
1930-32	2.307	5.390	3.161	2.105	1.130	1.054	1.247
1951	1.022	1.057	1.000	1.012	1.045	1.059	1.049
1952	.955	·95 0	-887	.939	·96 o	.950	.951
1953	.910	.901	.855	-888	.952	·9 5 4	·946
1954	∙868	•766	.742	·86 7	.945	.950	•936
1955	·857	.738	.790	·852	.945	.975	. 949
1956	·824	.702	.710	·821	.940	.971	.942
1957	∙806	·738	.774	·8 ₃₇	.951	.935	.924
1958	·776	.681	·726	·816	.933	·958	.929
1959	753	.709	.710	821	.929	·947	.921
1960	.772	.667	.742	.816	.913	924	.903
1961	.738	.723	.710	·816	.932	965	.932
Females							
1930-32	2.256	5.716	4.046	2.279	1.439	1.193	1.362
1951	1.008	1.083	.977	1.013	1.036	1.066	1.053
1952	·965	·88 ₃	∙886	·88 ₃	.951	·924	·9 2 6
1953	.929	.917	∙886	·851	.927	·926	.919
1954	·858	·69 2	.773	825	.919	·905	∙896
1955	857	.792	•796	.760	·9 o 6	.935	.913
1956	.813	•683	.705	.727	·891	·9 2 6	∙899
1957	·816	.717	.750	.747	.891	∙870	∙861
1958	.781	.625	•636	.708	·874	·897	.872
1959	.779	.675	.705	695	859	·888	·86 2
1960	.773	633	.705	·662	·843	·862	•839
1961	·746	·6 ₅ 8	•659	·669	·86 o	.900	∙869