346

THE RECENT TREND OF MORTALITY IN GREAT BRITAIN

By L. V. MARTIN, F.I.A.

of the Government Actuary's Department

In continuation of the tables given in *J.I.A.* **89**, 289, central death-rates in age-groups for males and females separately in the year 1963, and the ratios of those death-rates to the corresponding rates for the years 1950–52 taken as the standard, are shown in Table 1. The deaths are those recorded

Table 1.

Death rates for age-groups in Great Britain in the year 1963 and ratios of these rates to the rates for 1950–52

	Death-rate	s in 1963	Ratios to 1950-52 rates		
Age-group	Males	Females	Males	Females	
Under 1	.02504	·01940	.73	.73	
1-4	.00101	.00083	.72	.69	
5–9	·00050	.00032	·74	·67	
10–14	-00040	-00024	-73	·62	
15–19	.00088	.00035	.92	·51	
20-24	.00111	·00044	⋅80	·44	
25-29	.00101	·00062	·67	.50	
30-34	.00123	·00091	.70	.62	
35-39	.00188	.00142	·81	·74	
40-44	·00317	.00222	⋅89	·82	
45-49	.00553	.00369	.88	.88	
50-54	·00966	.00545	.88	·84	
55-59	·01736	.00849	.95	·86	
60-64	·03001	·01409	1.02	.88	
65-69	·0453	·0234	1.00	.88	
7074	·0712	·0405	1.03	.88	
75–79	·1089	·0705	.99	⋅88	
80-84	·168	·122	∙98	∙91	
85 and over	·284	·223	.99	.95	
All ages:					
Crude rates	·01289	-01159	1.013	1.039	
Standardized*	·01219	00967	.958	·867	

^{*} Standardized on 1951 Census age distribution

as occurring in Great Britain in the year and the population used is the 'home' population at mid-1963, i.e. the number of people actually in the country at the time, as estimated by the Registrars General of England and Wales and of Scotland.

The estimated population has been adjusted to take account of the results, so far as they are available, of the 1961 Census. Most of the adjustments are relatively small and do not materially affect comparisons with earlier years. One adjustment was, however, more substantial; the estimated number of women aged 85 and over has been increased

Table 2.

Standardized death-rates for age-groups in Great Britain (standardized on 1951 Census age distribution)

Year	Age-group						All	
	Under 1	1-4	5–14	15–44	45–64	65 and	ages	
						over		
Males								
1930–32	·07953	·00760	.00196	.00412	·01637	·0873	·01588	
1953	.03139	·00127	.00053	.00174	·01392	.0790	·01204	
1954	.02992	.00108	.00046	.00170	.01382	.0787	·01192	
1955	.02955	·00104	·00049	.00167	.01382	.0807	.01208	
1956	.02842	.00099	.00044	.00161	·01375	·0804	·01199	
1957	·02779	.00104	.00048	.00164	-01391	·0774	·01176	
1958	·02676	·00096	.00045	-00160	·01364	·0793	-01183	
1959	·02597	·00100	·00044	.00161	.01358	·0784	.01173	
1960	·02663	.00094	·00046	.00160	·01335	.0765	·01149	
1961	.02546	$\cdot 00102$	·00044	.00160	·01363	·0799	·01186	
1962	.02639	00092	·00045	·00156	·01364	.0802	·01188	
1963	.02504	·00097	.00045	·00158	·01395	.0828	·01219	
Females								
1930–32	∙05973	.00686	.00178	.00351	·01243	.0735	·01519	
1953	·02461	.00110	.00039	·00131	.00801	.0585	·01025	
1954	·02272	·00083	·00034	.00127	.00794	.0572	·00999	
1955	.02270	·00095	.00035	.00117	.00783	·0591	·01018	
1956	·02153	$\cdot 00082$.00031	.00112	·00770	·0585	·01002	
1957	.02160	·00086	·00033	·00115	-00770	·0550	-00960	
1958	-02068	·00075	·00028	·00109	·00755	·0567	·00972	
1959	·02062	·00081	·00031	-00107	·00742	-0561	·00961	
1960	-02048	·00076	.00031	·00102	·00728	.0545	∙00936	
1961	∙01976	·00079	.00029	·00103	·00743	∙0569	·00969	
1962	·019 7 0	·00077	.00030	·00104	.00739	.0560	.00957	
1963	.01940	.00080	·00028	·00103	.00749	∙0567	·00967	

by about 5%, resulting in a corresponding decrease in the apparent death-rate.

In order that the trend of mortality rates may be conveniently studied, the group death-rates and the ratios to the 1950-52 standard since 1953

are set out in Tables 2 and 3 and the ratios are shown graphically in Figs. 1 nd 2. Figures for 1952 and earlier years have not been given, but they are available in previous articles. In these tables and graphs the age-groups have been reduced to six—under age 1 (infants), 1–4 (young children), 5–14 (children of school age), 15–44 (younger adults), 45–64 (the middle

Table 3.

Ratios of standardized death-rates for age-groups in Great Britain to the rates for 1950–52 (standardized on 1951 Census age distribution)

Year	Age-group						All		
Year	Under 1	1–4	5–14	15-44	45-64	65 and over	ages		
Males									
1930–32	2.307	5.390	3.161	2.102	1.120	1.054	1.247		
1953	∙910	.901	⋅855	·888	.952	∙954	.946		
1954	·868	·766	·742	·867	.945	·950	·936		
1955	⋅857	·738	·790	⋅852	.945	.975	.949		
1956	·824	·702	·710	·821	·940	.971	·942		
1957	⋅806	·738	·774	·837	·951	.935	·924		
1958	·776	∙681	·726	·816	.933	·958	.929		
1959	·753	·709	.710	·821	.929	.947	.921		
1960	.772	·667	·742	∙816	.913	.924	.903		
1961	·738	·723	.710	·816	.932	.965	.932		
1962	·765	·652	·726	·796	.933	.969	.933		
1963	·726	·688	·726	⋅806	.954	1.000	.958		
Females									
1930–32	2.256	5.716	4.046	2.279	1.439	1.163	1.362		
1953	.929	·917	⋅886	·851	·927	.926	.919		
1954	⋅858	·692	.773	·825	.919	.905	∙896		
1955	⋅857	.792	·796	.760	·906	.935	.913		
1956	⋅813	·683	·705	·727	⋅891	·926	.899		
1957	∙816	·717	·750	.747	∙891	·870	∙861		
1958	·781	·625	·636	·708	⋅874	∙897	·872		
1959	·779	·675	·705	.695	⋅859	⋅888	·862		
1960	·773	·633	·705	.662	·843	·862	·839		
1961	·746	·658	.659	.669	.860	.900	·869		
1962	·744	·642	.682	·675	⋅855	⋅887	·858		
1963	·733	.667	·636	.669	⋅867	⋅897	⋅867		

aged) and 65 and over (the elderly). Further, the group death-rates and ratios have been standardized on the age-distribution in Great Britain at the 1951 Census so as to eliminate the effect of changes in the age-distribution of the population.

As a link with the earlier series of tables that appeared in the *Journal*, showing the trend of mortality in England and Wales since 1930–32, figures for Great Britain for the average of those three years have also been included in Tables 2 and 3.




Fig. 1. Ratios of standardized group death-rates to 1950-52 rates—children

The tables show that mortality in Great Britain in 1963 was heavy. For men, the standardized death-rate was about $2\frac{1}{2}\%$ greater than in 1962,

and was higher than in any year since 1951. For women, the increase over 1962 was only about 1%, but it would have been about $\frac{1}{2}\%$ more had it not




Fig. 2. Ratios of standardized group death-rates to 1950-52 rates—adults and all ages

been for the adjustment to the 85 and over population referred to above. No doubt the heavier mortality was largely brought about by the exceptionally severe weather in the early months of 1963.