



England & Wales mortality monitor – COVID-19 update – week 30 of 2020

Summary

There have been around 62,100 more deaths in the UK from the start of the pandemic to 24 July 2020 than if mortality rates were similar to those experienced in 2019. We have modified the method that we use to calculate the UK figure. We now use data from National Records Scotland (NRS) and the Northern Ireland Statistics and Research Agency (NISRA) as well as the Office for National Statistics (ONS), rather than scaling the England & Wales excess death figure using data from Public Health England (PHE) and the Department of Health and Social Care (DHSC). The new method increases the UK estimate by about 1,000.

There were 4% fewer deaths registered in England & Wales in week 30 of 2020 than would have been expected if standardised mortality rates had been the same as week 30 of 2019. Mortality was 4% lower in week 29 and 7% lower in week 28 than in the corresponding weeks of 2019.

The cumulative mortality improvement in England & Wales for 2020 is –10.6% as at 24 July 2020, compared to +0.1% as at 20 March 2020, before the coronavirus pandemic had a material impact.

Background

During the coronavirus pandemic we have been publishing frequent updates to the CMI Mortality Monitor. This update shows the position as at 24 July 2020 (week 30 of 2020), based on provisional deaths data published by the Office for National Statistics (ONS) on 4 August 2020. The previous monitor was for week 28, and we intend to publish the next, for week 32, on 18 August 2020. All updates are publicly available from the CMI pages of the Institute and Faculty of Actuaries website, together with software that we have made available to Authorised Users to carry out their own ad hoc analyses: <https://www.actuaries.org.uk/learn-and-develop/continuous-mortality-investigation/other-cmi-outputs/mortality-monitor>.

Notes

Full details of the methods used for results based on the ONS data are included in [Working Paper 111](#). Our analysis is based on Standardised Mortality Rates (SMRs). These adjust the provisional weekly deaths data published by the ONS to control for changes in the size, age and gender distribution of the population over time.

We have included versions of Charts D and E from the standard quarterly monitor, which show results for males and females combined, for ages 20-100:

- Chart 1 (like Chart D from the quarterly report) shows cumulative standardised mortality for each year, relative to the average for 2010-2019.
- Chart 2 (like Chart E from the quarterly report) shows cumulative standardised mortality improvements for each year (i.e. the progression of annual mortality improvements over the course of each year).
- Mortality rates and mortality improvements vary by age, and the results shown are sensitive to the age distribution of the chosen standard population (the 2013 European Standard Population).

Our calculations rely on data for registered deaths, and we are conscious that in recent weeks deaths may have been registered earlier or later than in previous years. Consequently, comparisons of mortality between 2020 and earlier years may not be on a like-for-like basis. Also, results for individual weeks may not be consistent between years due to the timing of public holidays.

Use of this document

The CMI disclaims any liability from use of or reliance on these calculations, including in relation to financial transactions such as longevity swaps; and the CMI does not guarantee that it will continue to publish updates. Please also see the reliances and limitations, disclaimer, and copyright notice on the final page of this document.

TAS compliance

This paper is intended to analyse recent mortality in England & Wales. It complies with the principles in the Financial Reporting Council's Technical Actuarial Standard "TAS 100: Principles for Technical Actuarial Work". Any person using this paper should exercise judgement over its suitability and relevance for their purpose.



Results

Chart 1 shows cumulative standardised mortality rates compared to the 2010-2019 average. Cumulative mortality to week 30 of 2020 is higher than cumulative mortality to week 30 in any year since 2007, and is 6.4% above the 2010-19 average. It was 1.9% below the 2010-19 average at week 12, before the coronavirus pandemic had a material impact. The highest value was +7.4% at week 23.

Chart 1: Cumulative standardised mortality rate compared to the 2010-2019 average





Chart 2 shows the cumulative annual standardised mortality improvement for 2020 and for the previous ten years. Note that Chart 2 shows cumulative improvements, so a higher value represents a higher improvement and lower mortality; whereas in Chart 1 a higher value represents higher mortality.

The cumulative mortality improvement is -10.6% as at 24 July 2020 (week 30 of 2020), compared to +0.1% as at week 12, before the coronavirus pandemic had a material impact. The lowest value was -11.1% as at week 24.

Chart 2: Cumulative annual standardised mortality improvement





Impact of coronavirus on total deaths

The ONS data shows 217 deaths registered in week 30 “where COVID-19 was mentioned on the death certificate”. The overall impact of the coronavirus pandemic on total deaths may be different:

- There may have been some deaths that were wholly or partially due to COVID-19 but where COVID-19 was not mentioned on the death certificate.
- Some deaths where COVID-19 was mentioned on the death certificate may not be “excess” deaths, as the deceased might have died from another cause in the same period, in the absence of the coronavirus.
- There may have been “forward mortality displacement”: some deaths that occurred earlier in the pandemic would otherwise have occurred in this week.
- There may have been indirect impacts on deaths due to restrictions on movement due to the coronavirus; for example, changes in traffic, pollution and mental health.

To consider the possible impact of the pandemic on total deaths, we have estimated the number of deaths that we would have seen in week 30 of 2020 if the SMRs for each gender and age-group had been the same in week 30 of 2020 as in week 30 of 2019. As mortality in the first 12 weeks of 2019 and 2020 was similar, as seen in Charts 1 and 2, this gives a broad indication of “expected” mortality in the absence of the coronavirus pandemic¹. We can then subtract the expected deaths from actual deaths to estimate the “excess” deaths that, in the absence of other likely causes, may be attributable to the pandemic.

We have not made any adjustment for differences in the timing of public holidays in 2019 and 2020. While such differences may affect expected, actual and excess results for individual weeks, positive and negative impacts for different weeks should cancel out over time in cumulative results.

Table 1 shows results for week 30, and for week 29, for which we did not produce a mortality monitor:

- Actual deaths in week 30 are 4% lower than expected: 3% lower than expected for males and 4% lower for females. In week 29 they were 4% lower than expected: 6% lower for males and 2% lower for females.
- Weeks 25 to 30 are the only weeks since late March that mortality in 2020 has been lower than in 2019.
- Although excess deaths are negative, there were 217 registered deaths reported by the ONS in week 30 where COVID-19 was mentioned on the death certificate.

Table 1: Comparison of COVID-19 deaths and “excess” deaths

Description	Week 30 of 2020			Week 29
	Male	Female	Total	Total
“Expected” registered deaths, if SMRs were the same in 2019 and 2020	4,629	4,607	9,235	9,204
Actual registered deaths, from all causes	4,489	4,402	8,891	8,823
“Excess” registered deaths (actual minus expected)	-140	-205	-344	-381
Registered deaths where COVID-19 was mentioned on the death certificate	114	103	217	295
Excess as a proportion of expected	-3%	-4%	-4%	-4%

¹ Our calculation of excess deaths depends on the historical period that we use to estimate expected deaths. If we had used the average standardised mortality rates for 2015-19 rather than only 2019 to calculate expected deaths, without allowing for mortality improvements, then this would have reduced the excess deaths by 262 (from -344 to -606) in week 30, and reduced the cumulative excess at week 30 (shown in Chart 4) from 56,379 to 49,216, a difference of 13%. We reiterate our preference for using 2019 to estimate expected deaths in the absence of a pandemic, as 2019 and 2020 had similar mortality experience for weeks 1 to 12.



Chart 3 compares “excess” registered deaths and registered deaths where COVID-19 was mentioned on the death certificate in each week since week 13. While there were some deaths in weeks 11 and 12 where COVID-19 was mentioned on the death certificate, the level of excess deaths in those weeks is small compared to typical weekly volatility in deaths, so cannot be reliably estimated. Chart 4 is similar to Chart 3, but plots the cumulative numbers of deaths, since week 13.

The number of excess deaths was much higher than the number of deaths where COVID-19 was mentioned on the death certificate in weeks 13 to 17, but this is not the case in later weeks. Excess deaths have been lower than mentions of COVID-19 in weeks 21 to 30, and negative in weeks 25 to 30.

Chart 3: Comparison of deaths registered in each week (see text for details)





Chart 4: Comparison of cumulative registered deaths (see text for details)



Charts 5 and 6 show excess deaths as a proportion of expected deaths by age band for each week. This has tended to be higher for older age bands throughout the pandemic. We do not show results for ages below 45 as the relatively low numbers of deaths at those ages means that estimates of expected deaths would be unreliable. Similarly, we no longer show a table of excess mortality by age band, as the figures have become more uncertain as the numbers of deaths have reduced.

Chart 5: Excess as a proportion of expected in each week – males (see text for details)





Chart 6: Excess as a proportion of expected in each week – females (see text for details)





Excess deaths for the United Kingdom

The rest of this report is based on data for England & Wales, published by the ONS. In this section we consider excess deaths for the United Kingdom as a whole.

We estimate that the numbers of excess deaths from the start of the pandemic to 24 July 2020 are:

- 56,600 for England & Wales²; and
- 62,100 for the United Kingdom.

In each case, the excess figure is calculated relative to the number of deaths expected if mortality rates were similar to those experienced in 2019.

We previously derived our UK estimate from our England & Wales estimate by using data published by Public Health England (PHE), for England, and the Department of Health and Social Care (DHSC), for the devolved administrations of Northern Ireland, Scotland and Wales. This data relates to deaths of people who have had a positive test result for the coronavirus confirmed by a Public Health or NHS laboratory.

Following concerns that PHE/DHSC figures may not have been recorded consistently across the UK, we have revised our method. We now work directly with weekly data for all-cause mortality from National Records Scotland (NRS) and the Northern Ireland Statistics and Research Agency (NISRA). The new method produces an estimate of UK excess deaths that is around 1,000 higher than our previous method.

We note that the week 30 figure for Northern Ireland is not yet available. However this is unlikely to materially affect our estimate as excess mortality in Northern Ireland in recent weeks has been small, between -44 and +34 in the past four weeks for which we have data.

Data sources

The provisional weekly deaths are available from:

ONE (England & Wales):

<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/datasets/weeklyprovisionalfiguresondeathsregisteredinenglandandwales>

NRS (Scotland): <https://www.nrscotland.gov.uk/covid19stats>

NISRA (Northern Ireland): https://www.nisra.gov.uk/sites/nisra.gov.uk/files/publications/Weekly_Deaths.xls

² This cumulative figure, since the start of the pandemic, is slightly different to that in footnote 1, which shows the cumulative figure since week 13.



Reliances and limitations

The purpose of the weekly mortality monitor is to provide regular updates on standardised mortality in England & Wales during the coronavirus pandemic, adjusting ONS data to allow for changes in the size and age of the population.

The CMI aims to produce high-quality outputs and takes considerable care to ensure that the mortality monitor and the accompanying spreadsheet of results are accurate. However:

- We cannot guarantee their accuracy (see the Disclaimer).
- There is a reliance on the data published by the ONS and described as provisional.
- We have also applied judgement and assumptions in deciding on the calculation methods and the presentation of results.
- Anyone using the results of the mortality monitor should ensure that it is appropriate for their particular use, and note that care is needed when estimating full year experience from partial year experience. This is particularly true during the coronavirus pandemic.

Disclaimer: This document has been prepared by and/or on behalf of Continuous Mortality Investigation Limited (CMI). The CMI does not accept any responsibility and/or liability whatsoever for the content or use of this document. Whilst care has been taken during the development of the document, CMI does not (i) warrant its accuracy; or (ii) guarantee any outcome or result from the application of this document or of any of CMI's work (whether contained in or arising from the application of this document or otherwise). You assume sole responsibility for your use of this document, and for any and all conclusions drawn from its use. CMI hereby excludes all warranties, representations, conditions and all other terms of any kind whatsoever implied by statute or common law in relation to this document, to the fullest extent permitted by applicable law. If you are in any doubt as to using anything produced by CMI, please seek independent advice.

Copyright: You may reproduce the contents of this document free of charge in any format or medium provided it is:

1. reproduced accurately and is unaltered;
2. not used in a misleading context; and
3. correctly referenced and includes both CMI's Disclaimer notice set out above and CMI's copyright notice, as follows:

© Continuous Mortality Investigation Limited.

Continuous Mortality Investigation Limited ("CMI") is registered in England & Wales
Company number: 8373631
Registered Office: 7th floor, Holborn Gate, 326-330 High Holborn, London, WC1V 7PP

Correspondence address: Two London Wall Place, 123 London Wall, London, EC2Y 5AU
Email: info@cmilimited.co.uk
Tel: 020 7776 3820

Website: www.cmilimited.co.uk (redirects to www.actuaries.org.uk)

Continuous Mortality Investigation Limited is wholly owned by the Institute and Faculty of Actuaries.